

As a mom, you
focus on
everyone else.

Now, it's your turn
to focus on you.



You are cordially invited to attend
My Breasts. My Body. *My Turn.*
our special event—just for moms



You do so much for your family. Now, make it your turn.
Come join other moms and get your questions
answered by a plastic surgeon.

To RSVP,

For more information, talk to your surgeon
and visit **ItsMyTurn.com.**

Nicole, age 36, mom of 1
Actual patient.
Individual results may vary.

Natrelle® Silicone-Filled Breast Implants
Important Information

Who may get breast implants (INDICATIONS)?

Natrelle® Silicone-Filled, Natrelle INSPIRA®, and Natrelle® 410 Shaped Gel Breast Implants are approved for women for the following:

- **Breast augmentation for women at least 22 years old for silicone-filled implants.**
Breast augmentation includes primary breast augmentation to increase breast size, as well as revision surgery to correct or improve the result of a primary breast augmentation surgery.

Please see Important Safety Information on next page.

IMPORTANT SAFETY INFORMATION

Who should NOT get breast implants (CONTRAINDICATIONS)?

Breast implant surgery should not be performed in women with active infection anywhere in their body, with existing cancer or pre-cancer of their breast who have not received adequate treatment for those conditions, and women who are currently pregnant or nursing.

What else should I consider (WARNINGS)?

Breast implants are not lifetime devices or necessarily a one-time surgery. You may experience unacceptable dimpling, puckering, wrinkling, or other cosmetic changes of the breast, which may be permanent. Breast implants may affect your ability to produce milk for breastfeeding. Silicone-filled implants may rupture without symptoms. You should have MRI examinations 3 years after surgery and then every 2 years thereafter in order to detect ruptures. Symptoms of a ruptured implant may be hard knots or lumps surrounding the implant or in the armpit, change or loss of size or shape of the breast or implant, pain, tingling, swelling, numbness, burning or hardening. Talk to your doctor about removing ruptured implants. Inform any other doctor if you have implants.

What types of conditions require more study (PRECAUTIONS)?

Caution: Notify your doctor if you have any of the following conditions, as the risks of breast implant surgery may be higher: autoimmune diseases, a weakened immune system, planned radiation therapy to the breast or planned chemotherapy following breast implant placement, conditions/medications that interfere with wound healing and blood clotting, reduced blood supply to breast tissue or a clinical diagnosis of depression, other mental health disorders, body dysmorphic disorder or eating disorders. These disorders should be resolved or stabilized prior to surgery.

What are some complications with breast implants (COMPLICATIONS)?

Key complications are reoperation, implant removal with or without replacement, implant rupture, and severe capsular contracture (severe scar tissue around the implant). Other complications include asymmetry, nipple/breast/skin sensation changes, scarring or wrinkling/rippling. Talk to your doctor about other complications.

Talk to your doctor. For more information see the Patient Brochures at www.allergan.com/labeling/usa.htm or call the Allergan Product Support line at 1-800-433-8871.

To report a problem with *Natrelle*® Silicone-Filled, *Natrelle* INSPIRA®, and *Natrelle*® 410 Shaped Gel Breast Implants, please call Allergan Product Surveillance at 1-800-624-4261.

Natrelle® Silicone-Filled, *Natrelle* INSPIRA®, and *Natrelle*® 410 Shaped Gel Breast Implants are available by prescription only.

