



## Lloyd Van Wyck, MD. FRCSC

1081 Carling Avenue, Suite 706  
Ottawa, Ontario K1Y 4G2  
(613) 722-7912

[www.cosmeticsurgeryottawa.com](http://www.cosmeticsurgeryottawa.com)

For two weeks prior to surgery and one week after surgery, do not take medications containing aspirin (ASA) (salicylates). Many over the counter medications contain aspirin. Please read the labels of any medications you are taking. If you are unsure whether or not your medication contains aspirin, please give the doctor or nurse the name of the medication. Please notify us of ANY and ALL medications you are currently taking, including over the counter medicines. You discontinue any herbal remedies and supplements.

### Medications to Avoid Prior to Surgery

Advil	Coumadin	Goody's	Sine Aid
Aleve	Darvon	Headache powder	Soma
Alka Seltzer	Dicumarol	and tablets	Tolectin
Alcohol	Disalcid	Ibuprofen	Toradol
Anacin	Dolobid	Indomethacin	Trandate
Anaprox	Dristan	Lodine	Trental
Ansaid	Easprin	Lovenox	Trilisate
Aspirin	Ecotrin	Meclomen	Vanquish
BC Tablets or powder	Empirin	Medipren	Vicoprofen
BC Cold powder	Equagesic	Midol	Vitamin E
Brufen	Enoxaparin	Monogesic	Voltaren
Bufferin	Excedrin	Motrin	Warfarin
Buffex	Feldene	Naprosyn	Wesprin
Cama arthritis pain reliever	Florinal	Pepto Bismol	Zavtrin
Celebrex	Flubiprofen	Percodan	Zoprin
Cephalgesics	Sodium	Persantine	
Cogesprin	4 Way cold tablets	Robaxisal	
Coricidin			

The diet drug known as Phen-Fed is marketed under the following generic and brand names. If you are on any of these drugs please inform this office. You must be off these drugs for at least two weeks prior to surgery or your surgery will be canceled.

PHENTERMINE	Phentride	Zantryl
Adipex-P	Phentride Caplets	Ephedra
Duromine	Phentrol	Ephedra
Fastin	Phentrol-2	Ephedrine
Oby-Mat	Phentrol-3	MaHuang
Oby-Nix	Phentrol-5	Meridia
Oby-Trim	T-Deit	Ginko
Panshape	Teramin	St. John's Wort
Phentercot	Wilpower	