

After Breast Augmentation Instructions

You will have a bandage/wrap in place after surgery. This should remain in place for the first 24 hours after surgery. You may remove your bandages the next morning after surgery.

It is very important that you **do not smoke, do not wear a nicotine patch, or do not chew nicotine gum** in the first 3 weeks after surgery. Severe scarring may result from failure to observe this precaution.

Medications (you may be given all or some of the following medications depending on your surgery):

- Pain Medication: You may take it every 4 hours as needed. Do not drive a vehicle or operate heavy machinery while using this medication.
- Anti-Nausea Medication: In the event that you experience nausea or vomiting, take this medicine. While you may not need it, it is advisable to fill the prescription just in case.
- Stool Softener: Your pain medicine can cause constipation, and excess strain and pressure can not only be unpleasant, but may hurt your incisions. Please take this while on the Pain Medication.
- Muscle Relaxant/Anti-anxiety Medication: This will help relax the muscles involved in your procedure, which will help with pain control and should help you sleep.

Although you have been given a prescription for pain medication, you may only need regular Tylenol, and should try it first. If your stomach becomes irritated or upset, stop the medication.

Do not use Ibuprofen or aspirin containing products which may increase your chances of bleeding complications.

Surgical site care:

- Leave bandages in place. They can be removed the morning after your surgery.
- After bandages are removed you may...
 - Remove gauze, shower gently and pat dry.
 - Do not allow direct high pressure shower spray to hit incisions.
 - Do not soak incisions.
 - Leave steristrips in place.
- Wear bra 23 hours a day for one month (to support the breasts).

Activities:

- Sleep with head of bed elevated 30 degrees.
- **No heavy lifting for two (2) weeks.** Nothing greater than 20lbs.
- Ambulate today/ tonight.
- Do not engage in sexual activity for a minimum of two weeks.

Returning to Particular Activities

- You may return to nonimpact aerobics at 2 weeks after surgery (stair master, low-speed elliptical).
- You may begin running at 4 weeks after surgery, but wear a sports bra over your surgical bra for the next 2 weeks (post-op week 5 and 6).
- You may begin push-ups, pull ups and swimming 6 weeks after your procedure.
- With all activities, it is important to listen to your body. Don't push too hard, take it "low and slow."
- Avoid quick, hard movements that could break a suture or tear the incision apart. Avoid excessive lifting that can cause increased pressure and possible bleeding.
- Avoid excessive downward pressure. Remember that your body is not used to the new weight of your breasts and we don't want the implant to move down past the incision line.

Diet:

- As tolerated, No restrictions.
- We would recommend you start resuming your normal diet gradually. Start with thin or thick liquids and advance to a general diet as tolerated.
- Drink plenty of water (eight 8-ounce glasses/day).
- Do not drink any alcoholic beverages for 3 days. Never drink alcoholic beverages while on prescribed pain medication.

Follow-up appointments: (Please call the Mercy Clinic Medical Spa at 417-820-9340 to schedule your appointment.)

- 1-3 days after surgery: Quick check after surgery.
- 2 weeks after surgery: Your steri strips will be removed at this time. After your steri strips are removed you may use regular moisturizer daily. Be sure to apply sunscreen to your suture lines to prevent pigmentation issues.
- 2 months after surgery: You will be instructed on scar care and breast massage at this visit.
- 6 months after surgery: Check incisions, breast implant location and for any residual swelling.
- Annually: We would love to make sure everything is going great after your breast augmentation procedure.

Watch for the following signs and symptoms. Notify Mercy Clinic Medical Spa (820-9340) immediately if these occur:

- Fever or blistering.
- Redness that looks like sunburn, tenderness or warmth in the area of redness.
- Sudden increase in one sided pain.
- Swelling that is much greater on one side.
- Bruising is **NORMAL**.

Call the Mercy Clinic Medical Spa 417-820-9340 if you have any questions.