

Cataract Medications Instructions	
Operative Date _____	Operative Eye _____
*Antibiotic Drops	*NSAIDS
<p>1 drop 3 times a day prior to surgery and continue for 7 days:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Besivance <input type="checkbox"/> Vigamox <input type="checkbox"/> Zymar <input type="checkbox"/> Ciprofloxacin <p>Different Instructions</p> <ul style="list-style-type: none"> <input type="checkbox"/> Poly Trim – 1 drop 4 times a day for 3 days prior continue for 7 days 	<ul style="list-style-type: none"> <input type="checkbox"/> Ilvero – 1 drop 1x a day 3 days prior and continue for 7 days <input type="checkbox"/> Bromday - 1 drop 1x a day 3 days prior and continue for 7 days <input type="checkbox"/> Prolensa - 1 drop 1x a day 3 days prior and continue for 7 days <input type="checkbox"/> Xibrom - 1 drop 2x a day 3 days prior and continue for 7 days <input type="checkbox"/> Nevanac - 1 drop 4x a day 3 days prior and continue for 7 days <input type="checkbox"/> Acular LS (Ketorolac) - 1 drop 4x a day 3 days prior and continue for 7 days <p>Diabetic patients must continue drops for 6 weeks post-operatively Right: _____ Date _____ Left: _____ Date _____</p>
*Steroids	Other Instructions
<ul style="list-style-type: none"> <input type="checkbox"/> PredForte – 1 drop 4x a day beginning day of surgery, then for 14 days, then decrease to 2x a day for more 14 days <input type="checkbox"/> Econopred - 1 drop 4x a day beginning day of surgery, then for 14 days, then decrease to 2x a day for more 14 days <input type="checkbox"/> Omnipred - 1 drop 4x a day beginning day of surgery, then for 14 days, then decrease to 2x a day for more 14 days <input type="checkbox"/> Prednisolone Acetate - 1 drop 4x a day beginning day of surgery, then for 14 days, then decrease to 2x a day for more 14 days <input type="checkbox"/> Durezol - 1 drop 2x a day beginning day of surgery, then for 14 days, then decrease to 1x a day for more 14 days <input type="checkbox"/> Lotemax - 1 drop 2x a day beginning day of surgery, then for 7 days, then decrease to 1x a day for more 7 days <p>*Give 5 minutes between instilling different medications</p>	<p>Same day or one day post-operative examination will be scheduled by your surgeon or by a doctor chosen by you or your surgeon. During the recovery process you will be placing drops in your eyes from 2-6 weeks depending upon your current eye health. Immediately following the procedure your vision is expected to be hazy due to the procedure. You must keep all follow-up appointments to reduce the risk of post-operative complications. Your eye maybe sore, red, scratchy, watery, tearing, or feel like you have a foreign body in it. You should refrain from strenuous activities for 7-10 days which will be determined by your surgeon. You may take Advil, Aleve, Motrin, or Tylenol for mild discomfort. You may also use a preservative-free artificial tears for eye dryness. You must avoid direct water flow into your face for 7 days when showering and avoid using hairdryers around any sutures. You will wear protective eye when sleeping for 2 weeks after surgery. For 1-2 days sleep 30 degrees elevate.</p> <p>If you experience an unprecedented reaction to the prescribed medication, you will need to contact our office: 850-266-7500</p> <p>We respectfully request that you contact our office if you experience any of the following symptoms: Fever, rash, discharge from the incision, swelling of the eyelids, unbearable light sensitivity, sudden decrease in vision, flashes of lights or floaters. Please call 850-266-7500 24 hours a day.</p> <p>Your next Appointment is: _____ Location: _____</p> <p>With Dr. _____</p>

***CRYSTALENS IOL PROCEDURES: PATIENT'S WILL RECEIVE AN ADDITIONAL DROP THAT WILL BLUR VISION FOR 3-5 DAYS**

All Additional questions should be referred back to the ME+FS patient's surgery packet